

# Synthesis of Bifunctional Monomers by the Palladium-Catalyzed Carbonylation of Cardanol and its Derivatives

## Authors

James E. Mgya,  
Dr. Stuart A. Bartlett,  
Dr. Egid B. Mubofu,  
Dr. Quintino A. Mgani,  
Prof. Alexandra M. Z. Slawin,  
Peter J. Pogorzelec,  
Prof. David J. Cole-Hamilton

○

## Abstract

A 1,2-bis(di-*tert*-butylphosphinomethyl)benzene-modified palladium catalyst has been used to synthesize bifunctional monomers of different chain lengths from cardanol. Short-chain derivatives of cardanol, such as (*E*)-3-(dodec-8-enyl)phenol; HOPhC<sub>12</sub>-ene, (*E*)-3-(undec-8-enyl)phenol; HOPhC<sub>11</sub>-ene, (*E*)-3-(dec-8-enyl)phenol; HOPhC<sub>10</sub>-ene, and 3-(non-8-enyl)phenol; HOPhC<sub>9</sub>-ene, were synthesized by the metathesis of cardanol with symmetrical internal alkenes. These derivatives were methoxycarbonylated to produce monomers with different chain lengths such as methyl-16-(3-hydroxyphenyl)hexadecanoate; HOPhC<sub>15</sub>COOMe, methyl-13-(3-hydroxyphenyl)tridecanoate; HOPhC<sub>12</sub>COOMe, methyl-12-(3-hydroxyphenyl)dodecanoate; HOPhC<sub>11</sub>COOMe, methyl-11-(3-hydroxyphenyl)undecanoate; HOPhC<sub>10</sub>COOMe, and methyl-10-(3-hydroxyphenyl)decanoate; HOPhC<sub>9</sub>COOMe, respectively. Polymerization of the synthesized monomers produced oligomers that consist of up to seven monomer units as confirmed by MALDI-TOF-MS. Lactone formation was also observed in some cases under polymerization conditions.