
Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

http://www.elsevier.com/authorsrights

Author's personal copy

Radiation Physics and Chemistry 91 (2013) 15–18

Contents lists available at SciVerse ScienceDirect

Radiation Physics and Chemistry

journal homepage: www.elsevier.com/locate/radphyschem

Determination of essential and toxic elements in clay soil commonly consumed by pregnant women in Tanzania

D. Mwalongo [a], N.K. Mohammed [b,*]

[a] Tanzania Atomic Energy Commission, PO Box 743, Arusha, Tanzania
[b] Department of Physics, University of Dar es Salaam, PO Box 35063, Dar es Salaam, Tanzania

HIGHLIGHTS

- We assessed exposure of heavy metals to pregnant mothers who consume geophagic soil.
- We analyzed 100 samples of soil originated in Tanzania.
- The technique used was energy dispersive X-ray fluorescent.
- Essential and toxic elements were detected in concentrations above WHO limits.
- Hence, geophagy is exposing pregnant mothers and their children to metal toxicity.

ARTICLE INFO

Article history:
Received 8 February 2013
Accepted 7 June 2013
Available online 14 June 2013

Keywords:
Essential elements
Toxic elements
Clay soil
Geophagy
Pregnant women

ABSTRACT

A habit of eating clay soil especially among pregnant women is a common practice in Tanzania. This practice known as geophagy might introduce toxic elements in the consumer's body to endanger the health of the mother and her child. Therefore it is very important to have information on the elemental composition of the eaten soil so as to assess the safety nature of the habit. In this study 100 samples of clay soil, which were reported to be originating from five regions in Tanzania and are consumed by pregnant women were analyzed to determine their levels of essential and toxic elements. The analysis was carried out using energy dispersive X-ray fluorescent technique (EDXRF) of Tanzania Atomic Energy Commission, Arusha. Essential elements Fe, Zn, Cu, Se and Mn and toxic elements As, Pb, Co, Ni, U and Th were detected in concentrations above WHO permissible limits in some of the samples. The results from this study show that the habit of eating soil is exposing the pregnant mothers and their children to metal toxicity which is detrimental to their health. Hence, further actions should be taken to discourage the habit of eating soil at all levels.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Geophagy is a human behavior common among pregnant women involving the ingestion of earthy substances such as soil and clay. Several studies agree that this behavior is rooted from evolution and is common among animal kingdom (Gilardi et al., 1999; Mahaney et al., 1996). The reasons for soil ingestion are not scientifically supported however; the habit is believed to be influenced by traditional, physiological, psychological, medicinal, cultural and religious beliefs (Hooda et al., 2004). For instance, in some parts of Africa, soil ingestion is considered to be medicinal for childbearing and for treatment of diarrhea and intestinal parasites (Hunter, 1973; Vermeer and Ferrell, 1985). Several studies have associated the cation exchange capacity (CEC) of the ingested soils with the properties of clay to control diarrhea which is attributed to the surface area and water retention capacity of the clay soil (Mahaney et al., 1996). Geophagy around the world can also be linked with deficiency of essential nutrients in human metabolism (Hooda et al., 2002). Iron (Fe) deficiencies which lead to anemia is thought to be a reason which attract pregnant and lactating woman to consume soil (Lanzkowsky, 1959; Danford, 1982). In Tanzania, this habit is practiced in many regions and common among lactating and pregnant woman. The clay soil which is made into a shape of a sausage is sold in markets across the country in places such as Kariakoo, Kilombero and Ngamiani in Dar es Salaam, Arusha and Tanga Regions, respectively.

Despite of all reasons that elucidate the motivation of soil eating practice, the concentration of metal consumed by geophagic individuals remain unknown. Therefore, quantification of essential and toxic elements from geophagy is important to assess the safety of the practice in the country.

2. Materials and methods

Hundred samples of soil sausages each of about 40 g were purchased at Kariakoo, Kilombero and Arusha main markets.

* Corresponding author. Tel.: +25 577 769 3229.
E-mail addresses: najat@udsm.ac.tz, njkassim@yahoo.com (N.K. Mohammed).

0969-806X/$ - see front matter © 2013 Elsevier Ltd. All rights reserved.
http://dx.doi.org/10.1016/j.radphyschem.2013.06.002

Author's personal copy

The samples which were reported to originate from Kigoma and Morogoro were purchased at Kariokoo market, those from Arusha and Kilimanjaro regions were bought at the main market in Arusha and those from Pemba were purchased in a market in Tanga region. The samples were identified by vendors as commonly eaten by pregnant women. Section 2.1 shows the regions where the soil samples are originating.

2.1. Map 1

In the laboratory, the soil was oven dried at 60 °C for 2 days until a constant weight was achieved. The dried samples were disaggregated with mortar and pestle and sieved through 0.5 mm polystyrene sieve and kept in well labeled polythene bags. A dry weight of 12 g of soil sample with 2.7 g of cellulose binder were thoroughly mixed, homogenized and be compressed into pellets of diameter of 32 mm to give reproducible irradiation and counting geometry. The elemental analyses of samples were conducted using a bench top energy dispersive X-ray spectrometer of TAEC in Arusha. The machine which is operated by automated turbo-quant X-lab ProTM software was operated at a rate of 50 W and 50 kV voltage and the florescent X-rays were collected by a Si(Li) detector having a resolution (FWHM) at Mn$K\alpha \leq 160$ eV. A spectrum run for 15 min gave a good continuity statistics and resolution of the peaks. The concentrations of individual elements were determined by using fundamental parameter method inbuilt in X-lab Pro computer software in which matrix effects was counted for. Quality control was carried out using IAEA soil 7, to verify the accuracy and reliability of the analytical data. Table 1 shows that the analyzed elements were within 13% accuracy.

3. Results and discussion

3.1. Minimum detection limit (MDL)

In this study the calculation of the MDL of each element was carried out using the turbo-quant method incorporated in the X-lab Pro software package. The following relation was used to determine the MDL values (Rousseau and Bouchard, 2005).

$$\mathrm{MDL} = \frac{3 \times C_i}{I_p - I_b} \sqrt{\frac{I_b}{T_b}} \quad (1)$$

where

T_b = Time used to measure background intensity
C_i = Concentration of the analyte
I_p = Peak intensity
I_b = Background intensity

The minimum detection limits of elemental concentrations in soil for EDXRF system used in this study are presented in Table 2.

3.2. Elemental concentrations in soil

The mean concentrations of essential elements and toxic heavy metals in clay soil originating from five regions in Tanzania are presented in Table 3.

3.2.1. Essential elements

As Table 3 shows, the ingested soils are a potential source of essential elements. The concentration of Fe was found to be higher in soils from Arusha and Kilimanjaro regions than soils from other regions. The lowest concentration of Fe was found in soil from Pemba. The Kilimanjaro and Arusha regions have high concentrations of Fe probably because their soils were developed from volcanic eruptions of several mountains in the regions. It is documented elsewhere that volcanic soil usually contains high concentration of Fe (Quantin, 2004; Garcia-Rodeja et al., 2004). Since this soil is eaten then it is important to compare the concentration of elements found in the soils with FAO/WHO permissible limits in food. The Fe concentrations found in the soils in this study were higher by the factor of 63, 459, 567, 752 and 699 for clay soil from Pemba, Kigoma, Morogoro, Arusha and Kilimanjaro, respectively, than the permissible level in food of 214 μg/g set by FAO/WHO (2001). Although the adequate iron intake is an important component in decreasing the incidence of anemia, high concentrations of Fe may lead to tissue damage, as a result of the formation of free radicals in the body (Gurzau et al., 2003).

The concentration of manganese (Mn) ranged from 120 ± 2.9 μg/g to 1215 ± 18 μg/g with lowest concentration for samples from Pemba and highest from Arusha. Manganese is an essential element which is present in metalloproteins, such as

Table 2
The minimum detection limit (MDL) in μg/g of the EDXRF system used in this study for elements in soil.

Elements	MDL	Elements	MDL
Ag	3.9	Mn	5.0
As	1.3	Ni	2.5
Bi	0.7	Pb	1.0
Cd	6.6	Se	0.6
Co	15.7	Sn	6.7
Cr	15.1	Th	1.0
Cu	2.1	U	2.8
Fe	5.0	V	31.5
Hg	1.4	Zn	1.1
I	18.5		

Table 1
Experimental and certified concentrations (μg/g ± SEM) of elements in reference materials IAEA Soil 7.

Element	Reference value(μg/g)	Experimental value	Element	Reference value (μg/g)	Experimental value(μg/g)
As	13.4 ± 0.9	14.1 ± 0.4	Mn	631 ± 4	647 ± 12
Al	47000 ± 21	43570 ± 40	Na	2400 ± 18	1700 ± 9
Ba	159 ± 9	155 ± 6	Ni	26 ± 3	26.5 ± 1.2
Br	7.0 ± 0.6	8.1 ± 0.5	P	460 ± 5	562 ± 12
Ca	163000 ± 62	161100 ± 231	Pb	60 ± 6	66 ± 1
Ce	61 ± 1	59 ± 2	Rb	51 ± 3	48 ± 2
Cr	60.0 ± 2.0	59.1 ± 0.6	Si	180000 ± 10	153400 ± 135
Cd	1.3 ± 0.1	1.2 ± 0.1	Sr	108 ± 4	103 ± 2
Cu	11.0 ± 0.7	12.6 ± 0.1	Th	8.2 ± 0.4	7.8 ± 0.9
Fe	25700 ± 85	26660 ± 44	Ti	3000 ± 13	3527 ± 28
Hg	0.04 ± 0.001	0.03 ± 0.1	U	2.6 ± 0.3	2.5 ± 0.7
K	12100 ± 45	11080 ± 14	V	66 ± 1	75 ± 2
Mg	11300 ± 39	7600 ± 22	Y	21 ± 0.6	24.5 ± 1.1

Author's personal copy

Table 3
The mean elemental concentrations (μg/g ± SEM) in soil with the maximum tolerable limits (MTLs) in food set by FAO/WHO 108.7 ± 7.5FAO/WHO (2001).

	Kigoma ($n=20$)	Morogoro ($n=20$)	Arusha ($n=20$)	Kilimanjaro ($n=20$)	Pemba ($n=20$)	MTL	Reference
As	9.9 ± 0.3	< 1.3	8.8 ± 2.3	< 1.3	< 1.3	0.05	WHO/IPCS (2002)
Cd	< 6.6	< 6.6	< 6.6	< 6.6	< 6.6	0.2	WHO (1992)
Co	55.9 ± 2.9	31.9 ± 1.3	32.0 ± 0.9	44.1 ± 8.7	23.4 ± 0.6	–	
Cr	71.7 ± 3.8	131.1 ± 4.5	207.6 ± 5.8	115.5 ± 6.9	423.2 ± 1.9	2.3	FAO/WHO (2001)
Cu	266.3 ± 5.8	55.7 ± 1.6	188 ± 11.2	33.1 ± 1.8	61.7 ± 3.1	40	FAO/WHO (2001)
Fe	98257 ± 131	121345 ± 212	160850 ± 132	149710 ± 143	13567 ± 98	214	FAO/WHO (2001)
Mn	377.8 ± 7.5	182.7 ± 1.7	1215.6 ± 18	162.5 ± 10.4	120.0 ± 2.9	1000	WHO (1999)
Ni	40.3 ± 0.4	29.4 ± 1.5	236.4 ± 4.9	31.7 ± 1.6	133.6 ± 8.4	0.2	FAO/WHO (2001)
Pb	12.1 ± 0.5	8.2 ± 0.3	32.3 ± 0.7	7 ± 0.4	14.6 ± 0.6	0.3	FAO/WHO (2001)
Se	1.2 ± 0.1	1.4 ± 0.1	< 0.62	1.2 ± 0.1	< 0.62		
Th	4.1 ± 0.1	2.4 ± 0.1	31.4 ± 2.1	2.4 ± 0.1	< 1.0	0.005	WHO (2001)
U	7.2 ± 0.2	< 2.8	18.5 ± 0.3	< 2.8	< 2.8	0.005	WHO (2001)
Zn	75.7 ± 2.1	54.6 ± 0.3	201.6 ± 11.1	57.9 ± 2.9	60		FAO/WHO (2001)

pyruvate carboxylase, and in the cytoplasmic glial enzyme, glutamine synthetase (WHO, 1999). It is reported that deficiency of manganese can cause severe skeletal and reproductive abnormalities in mammals while excess of manganese can lead to health effects primarily on the lungs and on the brain (WHO, 1999). WHO (1999) permissible limit for Mn in food is 1000 μg/g. The highest mean value of Mn in this study was found in samples from Arusha (1215) which is 1.2 times higher than the WHO permissible limit for Mn in food. The Mn concentration values found in samples from other regions were all within the permissible levels.

The concentration of copper (Cu) found in soils in this study ranged from 33.1 ± 1.8 μg/g to 266.3 ± 5.8 μg/g with the highest concentration found in clay soil from Kigoma and lowest in samples from Kilimanjaro. With the exception of clay soil from Kilimanjaro the concentration of Cu from Morogoro to Arusha were found to be higher than FAO/WHO permissible limits in food of 40 μg/g. The concentration of Zn varied from 54.6 ± 0.3 μg/g to 201.6 ± 11.1 μg/g with highest concentration in samples from Arusha and lowest in samples from Morogoro. The FAO/WHO permissible limit for Zn in foods is 60 μg/g, hence the values for Zn in soil samples from Morogoro, Kigoma and Pemba were above the recommended values (FAO/WHO, 2001).

Selenium (Se) was found in low concentrations in all soil samples analyzed in this study. Only three out of five analyzed regions had Se above the MDL for the EDXRF system. Chromium (Cr) which is also an essential element in trace amounts was found to vary from 71.7 ± 3.8 μg/g to 423.2 ± 1.9 μg/g. The Cr concentration is lowest for soil from Kigoma and highest for soil from Pemba. The FAO/WHO permissible limit for Cr in foods is 2.3 μg/g, hence the values for Cr in soil samples from all five regions were above the recommended values (FAO/WHO, 2001).

3.2.2. *Toxic heavy metals*

Nickel (Ni) is toxic even in low concentration as it may accumulate in the lungs and may cause bronchial hemorrhage or collapse (WHO, 2005). The concentration of Ni in the clay soil from five regions varied from 29.4 ± 1.5 μg/g to 236.4 ± 4.9 μg/g. Clay soil from Arusha region had the highest Ni concentration while Morogoro region had the lowest concentrations. The lowest value measured in these samples was 147 times higher than the permissible limit of 0.2 μg/g in food set by FAO/WHO (2001). Arsenic (As) was found in concentrations above MDL (1.3 μg/g) in samples from only two out of the five regions involved in this study. The value for the MDL of the EDXRF system used in this study is 26 times higher than the maximum permissible limit of 0.05 μg/g set by WHO/IPCS (2002)

Lead (Pb) is non-essential element as it is toxic even in trace amount (WHO, 2010). High concentrations in foodstuff have been reported to cause a number of diseases, including cardiovascular, renal failure (WHO, 1995). In this study the mean concentration of Pb varied from 7 ± 0.4 μg/g in samples from Kilimanjaro to 32.3 ± 0.7 μg/g, in samples from Arusha. All of these values are higher than the maximum permissible limit of 3 μg/g in food set by FAO/WHO (2001).

Cobalt (Co) was found in samples from all five regions studied in this work. The highest concentration was found in samples from Kigoma (55.9 ± 2.9 μg/g) and the lowest concentration was found in samples from Pemba (23.4 ± 0.6 μg/g). Cobalt has both beneficial and harmful effects on human health (WHO, 2006). It is beneficial for humans because it is part of vitamin B12, which is essential to maintain human health. High concentrations of Co have been reported to cause cancer. However, there are no suitable data with which to derive a tolerable intake for chronic ingestion of cobalt (WHO, 2006).

Uranium (U) and thorium (Th) were detected in values above MDL in samples from only two regions: Arusha and Kilimanjaro. These regions, both in northeastern Tanzania, have volcanic soil. The concentrations of U and Th were found highest in Arusha region with the concentration of 18.5 μg/g and 31.4 μg/g, respectively. Uranium and Th in food should be in very low concentrations. WHO set a permissible limit of 0.005 μg/g (WHO, 2001).

All of the measurements for the toxic element cadmium (Cd) were below the detectable limits for the EDXRF system MDL of 6.6 ug/g. However the WHO permissible limit for Cd in food is 0.2 μg/g, which is 33 times lower than the MDL of Cd in this study (WHO, 1992).

4. Conclusion

This study has determined concentration of essential and toxic elements in geophagic clays from five regions in Tanzania using energy dispersive X-ray fluorescent technique. The concentrations of five out of six essential elements were above the WHO permissible limits for these elements in consumed food. Although these elements are essential in the body activities, high concentrations may pose health problems. Toxic elements Pb, Ni and Co were detected in concentrations above WHO permissible limits in all samples analyzed in this study. U and Th which are toxic as well as radioactive elements were detected in samples from Arusha and Kilimanjaro which are Volcanic Regions. Arsenic was found in concentrations above MDL value of 0.4 μg/g in samples from two out of five regions involved in this study. Cd was not found in any of the samples in concentrations above MDL. However, the MDL of

Author's personal copy

the system used in this study was 6.6 μg/g which is 33 times higher than the WHO permissible limit of 0.2 μg/g. As Cd is toxic and carcinogenic further analysis using instrument with lower MDL is recommended. The results from this study show that the habit of eating soil is exposing the pregnant mothers and their children to metal toxicity which is detrimental to their health and the health of their children. Hence, the study is recommending that the habit of eating soil in Tanzania should be discouraged.

Acknowledgements

The Authors would like to acknowledge the staff of Tanzania Atomic Energy Commission in Arusha for their technical advice and support.

References

Danford, D.E., 1982. Pica and nutrition. Ann. Rev. Nutr 2, 303–322.
FAO/WHO, 2001. Food Additives and Contaminants, Joint Codex Alimentarius Commission, FAO/WHO. Food standards Programme, ALINORM 01/12A.
Garcia-Rodeja, E., Novoa, J.C., Pontevedra, X., Martinez-Cortizas, A., Buurman, P., 2004. Aluminium fractionation of European volcanic soils by selective dissolution techniques. Catena 56, 155–183.
Gilardi, J.D., Duffey, S.S., Munn, O.A., Tell, L.A., 1999. Biochemical functions of geophagy in parrots: detoxification of dietary toxins and cytoprotective effects. J. Chem. Ecol. 25, 897–922.
Gurzau, E.S., Neagu, C., Gurzau, A.E., 2003. Essential metals—case study on iron. Ecotoxicol. Environ. Saf 56, 190–200.
Hooda, P.S., Henry, C.J.K., Seyoum, T.A., Armstrong, L.D.M., Fowler, M.B., 2002. The potential impact of geophagia on the bioavailability of iron, zinc and calcium in human nutrition. Environ. Geochem. Health. 24, 305–319.
Hooda, P.S., Henry, C.J.K., Seyoum, T.A., Armstrong, L.D.M., Fowler, M.B., 2004. The potential impact of soil ingestion on human mineral nutrition. Sci. Total Environ. 333, 75–88.
Hunter, J.M., 1973. Geophagy in Africa and the United States: a culture-nutrition hypothesis. Geogr. Rev. 63, 173–195.
Lanzkowsky, P., 1959. Investigation into the etiology and treatment of Pica. Arch. Dis. Childhood 34, 140–148.
Mahaney, W.C., Hancock, R.G.V., Aufreiter, S., Huffman, M.A., 1996. Geochemistry and clay mineralogy of termite mound soil and the role of geophagy in chimpanzees of the Mahale Mountains, Tanzania. Primates 37, 121–143.
Quantin, P., 2004. Volcanic soils of France. Catena 56, 95–109.
Rousseau, R.M, Bouchard, M., 2005. Fundamental algorithm between concentration and intensity in XRF analysis—2, practical application. J. Anal. Chem., X-ray Spectrom. 13, 121–125.
Vermeer, D.E., Ferrell Jr., R.E., 1985. Nigerian geophagical clay: a traditional anti-diarrheal pharmaceutical. Science 227, 634–636.
WHO, 1992. Cadmium. Environmental Health Criteria. World Health Organization, Geneva p. 280.
WHO, 1995. Lead. Environmental Health Criteria. World Health Organization, Geneva p. 65.
WHO, 1999. Manganese and its Compounds. World Health Organization, Geneva.
WHO, 2001. Depleted Uranium: Sources, Exposure and Health Effects. Department of Protection of the Human Environment, Geneva.
WHO/IPCS, 2002. Arsine: Human Health Aspects. World Health Organization, International Programme on Chemical Safety (Concise International Chemical Assessment Document No. 47), Geneva.
WHO, 2005. Nickel in Drinking-water. Background Document for Development of WHO Guidelines for Drinking-water Quality. World Health Organization, Geneva. (WHO/SDE/WSH/05.08/55).
WHO, 2006. Cobalt and Inorganic Cobalt Compounds. World Health Organisation, Geneva.
WHO, 2010. Childhood Lead Poisoning. World Health Organization, Geneva.